



DOSSIER · SUBSTANCE

Biotin (Vitamin B7)

Biotin is a water-soluble B vitamin; human studies found lower total cholesterol in type 2 diabetes and mixed findings for hair, skin, nails and multiple sclerosis.

413

STUDIES IN
TOTAL

161

IN HUMANS

30

OF WHICH
CONTROLLED

30

IN ANIMALS

29

IN THE LAB

Figures from the IHR study database · Last updated 2026-10-17

How to Read This Dossier

ESTABLISHED

measured in humans, with a comparison group

PLAUSIBLE

supported by mechanism, animal studies or observation

HYPOTHESIS

a likely connection that still needs to be tested

Biotin (Vitamin B7) at a Glance

In type 2 diabetes, total cholesterol fell with biotin in a meta-analysis; the reduction in fasting blood glucose and triglycerides was not statistically confirmed, insulin remained unchanged ².

In hair loss, a systematic review found no difference in hair growth between biotin and placebo in the most informative, placebo-controlled study ⁴.

In progressive multiple sclerosis, disability regressed in some patients with high-dose biotin in a randomized study, and in no one with placebo ⁷. The follow-up study SPI2 did not confirm this benefit ²⁵.

Another randomized controlled study showed no difference in visual acuity, nerve fiber layer thickness and quality of life in MS-related chronic vision loss ⁹.

We evaluated 123 human studies on Biotin (Vitamin B7), including 12 randomised controlled trials and 6 meta-analyses or systematic reviews; the controlled studies include 1,170 participants in total. As of 11 October 2026.

1. What Is Biotin (Vitamin B7)? Occurrence, Origin and Forms

Biotin is a water-soluble vitamin of the B group. In the body it acts as a coenzyme of several carboxylases involved in fat, carbohydrate and protein metabolism. As a supplement it is traditionally associated with hair, skin and nails, and medically it is given in confirmed biotin deficiency and in inborn disorders of biotin metabolism.

Biotin occurs in many foods, among them egg yolk, liver, nuts, oats, soy and yeast, and is additionally produced by gut bacteria. In body cells it is present bound to protein.

In foods biotin is bound to protein and is released in the gut by the enzyme biotinidase. In supplements it is present in free form; human studies examine free biotin taken by mouth, high-dose biotin and biotin as a component of combination products, and in metabolic medicine also biotin as an infusion.

Forms and Preparations Studied

- Biotin from foods, bound to protein
- Free biotin as a dietary supplement (tablets, capsules, drops)
- High-dose biotin as used in studies of neurological disease
- Biotin in combination products with other vitamins and minerals
- Biotin for injection or infusion in inborn metabolic disorders

Not covered on this page:

- Other B vitamins such as thiamine, riboflavin, pantothenic acid, folate and cobalamin are substances in their own right, even when they occur in the same products.
- Combination products for hair, skin and nails that contain zinc, selenium or amino acids alongside biotin are not studies of biotin alone.
- Interference of laboratory tests by biotin, for instance in immunoassays, is an analytical safety topic and not an area of application.

2. What Does Biotin (Vitamin B7) Do? Results by Area

AREA	RESULT IN CONTROLLED STUDIES	STRENGTH OF EVIDENCE	STUDIES	PARTICIPANTS
Biotin deficiency	mostly improved	moderate	0 meta-analyses · 3 controlled · 24 other	128
Multiple sclerosis	mostly no difference	low	2 meta-analyses · 2 controlled · 20 other	247
Hair loss	mixed	unclear	1 meta-analyses · 2 controlled · 9 other	165
Skin and skin inflammation	mostly no difference	unclear	0 meta-analyses · 2 controlled · 5 other	165
Glucose and lipid metabolism	mixed	unclear	1 meta-analyses · 1 controlled · 5 other	40
Brittle nails	mixed	unclear	0 meta-analyses · 1 controlled · 2 other	105
Other investigations	mostly no difference	unclear	0 meta-analyses · 2 controlled · 51 other	146

How to read the table: counted are human studies whose results concern the area. “High” means at least one meta-analysis and two controlled studies showing improvement, or at least three controlled studies of which three quarters show improvement, with at least 300 participants in total. “Moderate” means at least two controlled studies showing improvement, as a majority, or one meta-analysis alongside controlled studies. “Low” means one controlled study showing improvement. Where an area has no controlled study, the strength of evidence is at most “low” (result “reviews only” or “observational data

only"). Everything else is "unclear". A study counts as showing improvement when at least half of its outcomes in this area improved. Reviews of combination products do not count as meta-analyses; results that only describe side effects or shares of users do not count as an effect. The same study is counted only once.

3. What the Studies Show in Detail

Biotin (Vitamin B7) and Biotin Deficiency: Human Studies

In pregnant women with increased excretion of 3-hydroxyisovaleric acid, a sign of mild biotin deficiency, this decreased with biotin in a randomized trial, while it increased slightly with placebo ¹⁵. In children with severe protein-energy malnutrition, the activity of biotin-dependent enzymes in lymphocytes increased with biotin, most strongly in those with initially low levels ²⁰. The evidence is considered moderate, as both studies are small and involve different populations and outcome measures.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
¹⁰ Hanson et al. 2026	randomised controlled · 80 participants	Prenatal multivitamin with 150 µg (intervention) or 30 µg (control) biotin; 150 µg vs. 30 µg biotin daily; 24 weeks;
¹⁵ Mock et al. 2002	randomised controlled · 26 participants	Biotin supplement or placebo; 300 µg (1.2 µmol) biotin per day; 14 days; vs. Placebo-controlled, randomized
²⁰ Velázquez et al. 1995	controlled · 22 participants	Biotin supplement; 10 mg/d; vs. Placebo, double-blind
²² Jezela-Stanek et al. 2022	cross-sectional study · 1,071,463 participants	Biotinidase activity in dried blood spot and serum, BTD gene sequencing
²⁶ Upadia et al. 2025	cohort study · 478 participants	Newborn screening for inborn errors of metabolism, including biotinidase deficiency; January 2005 to December
²⁷ Sharma et al. 2024	cross-sectional study · 407 participants	Biotinidase enzyme analysis and BTD gene sequencing; vs. enzyme activity <10% vs. >10% of mean normal
²⁸ Yaralı et al. 2024	cohort study · 357 participants	Biotin; vs. no control group
²⁹ Çıktı et al. 2024	cross-sectional study · 302 participants	Biotin (endogenous, in biotinidase deficiency); vs. genotype-phenotype correlation, no intervention comparison

Further studies in this area: ^{30 35 41 45 49 50 52 61 62 65 77 79 83 84 86 91 99 103 107}

Biotin (Vitamin B7) and Multiple Sclerosis: Human Studies

In multiple sclerosis, high-dose biotin (MD1003) has been tested against placebo in several randomized trials ⁷⁹²⁵. In a randomized trial in progressive MS, disability improved in some patients on high-dose biotin, and in none on placebo ⁷. EDSS progression was also lower ⁷. The follow-up study SPI2 did not confirm this benefit ²⁵. Another randomized controlled trial showed no difference in visual acuity, retinal nerve fiber layer thickness, and quality of life in MS-related chronic vision loss ⁹. A systematic review with meta-analysis found no confirmed difference from placebo in disability and EDSS, but more often a faster walking time ¹. The certainty of evidence is low: the first study showed benefit in some patients, but the follow-up study and the meta-analysis did not confirm this for disability. It remains open whether a subgroup benefits, for example in walking ability.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
¹ Espiritu et al. 2021	meta-analysis · 889 participants	High-dose biotin (MD1003), oral, as adjunctive treatment; at least 300 mg/day; 12 to 15 months; vs. Placebo
⁵ Marx et al. 2020	systematic review	Biotin as nutraceutical; vs. Control
⁷ Tourbah et al. 2016	randomised controlled · 154 participants	MD1003 (high-dose biotin); 100 mg biotin thrice daily; 12 months double-blind, then 12 months open-label for all; vs.
⁹ Tourbah et al. 2018	randomised controlled · 93 participants	MD1003 (high-dose pharmaceutical-grade biotin); 300 mg/day orally; 6 months double-blind, then 6 months open-la

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
²⁵ Joseph et al. 2026	study · 506 participants	High-dose biotin vs. placebo (SPI2 trial), arms pooled; Follow-up over 27 months; vs. Pooled treatment arms, no effi
³¹ Pape et al. 2020	cross-sectional study · 186 participants	vitamins and dietary supplements, including biotin; biotin 10 and 300 mg/day in healthy volunteers; vs. biotin intake v
³² Couloume et al. 2020	cohort study · 178 participants	High-dose biotin; 12 months; vs. No control group, comparison to baseline
³⁷ Collongues et al. 2021	cohort study · 48 participants	High-dose pharmaceutical-grade biotin (MD1003); 1 year; vs. No control group, before/after observation

Further studies in this area: [38](#) [46](#) [80](#) [81](#) [94](#) [95](#) [99](#) [100](#) [104](#) [110](#) [111](#) [112](#) [114](#) [115](#) [116](#) [117](#)

Biotin (Vitamin B7) and Hair Loss: Human Studies

A systematic review found three studies on biotin in hair loss; the most informative, placebo-controlled one showed no difference in hair growth ⁴. With biotin from a plant extract, with or without silica, hair loss decreased in a placebo-controlled study compared with baseline ⁸. A hair product for application with Redensyl, saw palmetto and biotin performed better together with platelet-rich plasma than Procapil with platelet-rich plasma ¹². Injections with biotin and dexpanthenol from two manufacturers reduced hair loss in both groups; there was no group without biotin ¹³. In acne patients taking isotretinoin, the proportion of growing hairs increased with additional biotin compared with baseline ¹⁸. It remains open whether biotin helps in hair loss mainly in a biotin deficiency.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
⁴ Yelich et al. 2024	systematic review	Oral biotin as a hair supplement; vs. Placebo or control groups in the included studies
⁸ Patel et al. 2025	randomised controlled · 105 participants	Botanical extracts with biotin from Sesbania grandiflora, some with silica from Bambusa arundinacea; 1.25 mg biotin or
¹⁸ Aksac et al. 2021	controlled · 60 participants	Biotin 10 mg/day added to isotretinoin; 10 mg/day; 4 months; vs. isotretinoin alone (group A) vs. isotretinoin plus biotin
²¹ Schulpis et al. 2001	case series	Biotin supplementation; 10 mg/day; 45 days of valproic acid treatment; vs. Healthy controls; three valproic acid dose gr
³³ Şen et al. 2021	cohort study · 156 participants	Biotin 1000 mcg/day; 1000 mcg/day; 3 months (biotin deficiency) or mean 2.5 months (self-initiated); vs. patients with ·
³⁴ Kakunje et al. 2025	cross-sectional study · 150 participants	Valproate treatment; measurement of serum biotinidase and IL-2; 2-year study period; vs. valproate group vs. control g
⁴³ Starace et al. 2021	case series · 25 participants	biotin alone or with topical minoxidil; vs. no control group, retrospective case series
⁴⁴ González et al. 2025	study · 22 participants	Serum WS Biotin, water-soluble form of D-biotin; 3 months daily application; vs. no control group (single-arm, pretest-i

Further studies in this area: [87](#) [96](#) [102](#) [109](#)

Biotin (Vitamin B7), Skin and Skin Inflammation: Human Studies

In acne patients taking isotretinoin, skin moisture decreased significantly without biotin, but not with additional biotin; a direct group comparison was not reported ¹⁸. With biotin from a plant extract, with or without silica, skin elasticity increased from baseline in a placebo-controlled study ⁸. Both studies are small, one was not randomized, and neither reports a direct comparison with the control group. It remains open whether the effects also occur in individuals with adequate biotin status without concomitant medications.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
⁸ Patel et al. 2025	randomised controlled · 105 participants	Botanical extracts with biotin from Sesbania grandiflora, some with silica from Bambusa arundinacea; 1.25 mg biotin
¹⁸ Aksac et al. 2021	controlled · 60 participants	Biotin 10 mg/day added to isotretinoin; 10 mg/day; 4 months; vs. isotretinoin alone (group A) vs. isotretinoin plus bic
²¹ Schulpis et al. 2001	case series	Biotin supplementation; 10 mg/day; 45 days of valproic acid treatment; vs. Healthy controls; three valproic acid dos
⁵² Ting et al. 2025	case series · 5 participants	Biotin supplementation; 10-30 mg daily; vs. no control group
⁵⁹ Pi et al. 2025	case series · 2 participants	Oral multivitamin regimen with biotin, lipoic acid, and pantothenic acid; vs. No control group; before-and-after cour
⁸⁶ Yilmaz et al. 2024	cross-sectional study	Biotinidase enzyme activity and BTD gene analysis; vs. genotype-phenotype correlations, no treatment groups
¹⁰⁸ Januszewski et al. 2023	review	Biotin among 13 other nutritional supplements

Biotin (Vitamin B7), Glucose and Lipid Metabolism: Human Studies

In adults with type 2 diabetes, total cholesterol fell with biotin in a meta-analysis; the reduction in fasting blood glucose and triglycerides was not statistically confirmed, insulin remained unchanged ². For mixtures of biotin and chromium picolinate, two randomized controlled studies reported improvements in blood glucose and lipid values in type 2 diabetes, but these findings apply to the mixture and not to biotin alone ⁶¹⁴. In the meta-analysis, only the reduction in total cholesterol is confirmed; whether biotin alone lowers blood glucose must be clarified by larger studies.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON	RESULT
² Zhang et al. 2022	meta-analysis · 445 participants	Biotin supplementation; 28 to 90 days; vs. Control groups of included RCTs	Fasting blood gluc
³ Chai et al. 2025	systematic review	Water-soluble vitamins, including biotin	Glycemic control a
¹⁹ Marshall et al. 1980	controlled · 40 participants	Biotin supplementation; 0.9 mg/day; 71 days; vs. Placebo	Total plasma lipids:
²⁴ Belda et al. 2022	study · 1,545 participants	Biotin plus fructo-oligosaccharides (prebiotics); vs. high-fat diet-fed mice without supplementation	weight gain and gly
⁶⁶ Walimbe et al. 2025	case report · 1 participants	Targeted replacement with biotin, lipoic acid and pantothenic acid	Metabolic control:
⁹² Güder et al. 2023	cross-sectional study	biotin (level measurement, no supplementation); vs. no control group, correlations	biotin levels and cr
⁹⁸ Mehta et al. 2026	review	Biotin and/or chromium supplements; vs. Single versus combined supplementation, controls (qualitative)	Fasting blood gluc

Biotin (Vitamin B7) and Brittle Nails: Human Studies

With biotin from a plant extract, with or without silica, nail roughness decreased from baseline in a placebo-controlled study with 105 participants ⁸. A review names biotin alongside collagen peptides, keratin, MSM, and orthosilicic acid among substances that may improve the appearance, strength, and brittleness of nails ¹⁰⁵. The certainty of evidence is unclear because only one controlled study is available, which does not report a comparison with placebo, and it remains open whether the participants had a biotin deficiency.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
⁸ Patel et al. 2025	randomised controlled · 105 participants	Botanical extracts with biotin from Sesbania grandiflora, some with silica from Bambusa arundinacea; 1.25 mg biotin
⁹⁷ Fernández-Campos et al. 2020	study	Regenail® nail lacquer with biotin, MSM, dimethylsilanediol salicylate; vs. Betalfatrus® reference product
¹⁰⁵ Zaraa et al. 2025	review	Biotin (dietary supplement)

Biotin (Vitamin B7): Further Outcomes in Studies

In children with asthma, biotin in addition to standard therapy reduced the inflammatory markers TNF-α and 8-OHdG more than placebo; lung function improved only with L-carnitine ¹¹. The other works in this area are predominantly observational studies or investigate biotin in combination products; controlled studies on clinical endpoints are still pending here.

STUDY	DESIGN · PARTICIPANTS	PREPARATION · DOSE · DURATION · COMPARISON
¹ Espiritu et al. 2021	meta-analysis · 889 participants	High-dose biotin (MD1003), oral, as adjunctive treatment; at least 300 mg/day; 12 to 15 months; vs. Placebo
¹⁰ Hanson et al. 2026	randomised controlled · 80 participants	Prenatal multivitamin with 150 µg (intervention) or 30 µg (control) biotin; 150 µg vs. 30 µg biotin daily; 24 weeks; vs. Lc
¹¹ Khedr et al. 2026	randomised controlled · 66 participants	Biotin 5 mg capsules twice daily plus standard therapy; 5 mg twice daily; 3 months; vs. Placebo plus standard therapy
²³ Kong et al. 2025	cohort study · 144,439 participants	Dietary biotin intake (24-h Oxford WebQ); median 14.06 years follow-up; vs. Q1 (lowest intake) vs. Q2-Q4
²⁴ Belda et al. 2022	study · 1,545 participants	Biotin plus fructo-oligosaccharides (prebiotics); vs. high-fat diet-fed mice without supplementation
²⁶ Upadia et al. 2025	cohort study · 478 participants	Newborn screening for inborn errors of metabolism, including biotinidase deficiency; January 2005 to December 2024
³⁶ Jaafar et al. 2025	cross-sectional study · 50 participants	Biotin (serum and follicular fluid levels, no supplementation); vs. Normal, poor, and hyper-responders
³⁹ Saudubray et al. 1984	case series · 41 participants	Vitamin mixture with biotin, thiamine, B12, riboflavin; biotin 10 mg; vs. comparison with similar reported cases

Further studies in this area: [40](#) [42](#) [47](#) [48](#) [51](#) [52](#) [53](#) [54](#) [55](#) [56](#) [57](#) [58](#) [59](#) [60](#) [63](#) [64](#) [66](#) [67](#) [68](#) [69](#) [70](#) [71](#) [72](#) [73](#) [74](#) [75](#) [76](#) [77](#) [78](#) [82](#) [85](#) [86](#) [88](#) [89](#) [90](#) [91](#) [92](#) [93](#) [99](#) [100](#) [101](#) [106](#) [107](#) [113](#) [118](#)

4. How Is Biotin (Vitamin B7) Dosed? Preparation, Dose and Duration in Studies

The studies mostly used oral biotin: as a dietary supplement, high-dose as a drug (MD1003) in multiple sclerosis or in combination products with other vitamins, minerals or plant extracts. Biotin was injected together with dexpanthenol in hair loss ¹³, applied in a hair product with Redensyl and saw palmetto ¹². The doses and dosage forms are listed in the table.

PREPARATION OR FORM	DOSE	DURATION	STUDY
Chromium picolinate plus biotin (CPB)	600 µg chromium and 2 mg biotin once daily	90 days	⁶
MD1003 (high-dose biotin)	100 mg biotin thrice daily	12 months double-blind, then 12 months open-label for all	⁷
Botanical extracts with biotin from Sesbania grandiflora, some with silica from Bambusa arundinacea	1.25 mg biotin orally; biotin-silica group additionally 21.75 mg silica	90 days	⁸
MD1003 (high-dose pharmaceutical-grade biotin)	300 mg/day orally	6 months double-blind, then 6 months open-label	⁹
Prenatal multivitamin with 150 µg (intervention) or 30 µg (control) biotin	150 µg vs. 30 µg biotin daily	24 weeks	¹⁰

PREPARATION OR FORM	DOSE	DURATION	STUDY
Biotin 5 mg capsules twice daily plus standard therapy	5 mg twice daily	3 months	11
Topical redensyl, saw palmetto, and biotin plus PRP	—	four sessions at three-week intervals	12
Weekly injections of biotin 5 mg/1 ml plus dexpantenol 250 mg/2 ml	Biotin 5 mg plus dexpantenol 250 mg weekly	6 weeks treatment, follow-up at 1 and 8 weeks	13
Chromium picolinate and biotin (CPB) or placebo	—	4 weeks	14
Biotin supplement or placebo	300 µg (1.2 µmol) biotin per day	14 days	15
SPP1 specialized food product with biotin and other vitamins	2 portions of SPP1 per day	14 days	16
AG1, comprehensive nutritional supplement with vitamins, minerals, probiotics, phytochemicals	Single serving 13 g mixed in water	Single dose, 1-week washout	17
Biotin 10 mg/day added to isotretinoin	10 mg/day	4 months	18
Biotin supplementation	0.9 mg/day	71 days	19
Biotin supplement	10 mg/d	—	20

5. Is Biotin (Vitamin B7) Safe? Side Effects in Studies

Adverse events with high-dose biotin in multiple sclerosis occurred no more frequently than with placebo [179](#); in children with asthma, biotin proved to be safe [11](#). However, high-dose biotin distorted laboratory tests in some of those treated, and in no one with placebo [1](#).

STUDY	REPORTED SIDE EFFECTS
1 Espiritu et al. 2021	Any and serious AEs not different; laboratory test interference in 4.7% on biotin
6 Albarracin et al. 2007	well tolerated, no adverse effects, similar to placebo
7 Tourbah et al. 2016	Safety profile similar to placebo
8 Patel et al. 2025	No adverse events reported
9 Tourbah et al. 2018	Safety profile similar to placebo
11 Khedr et al. 2026	Biotin was safe
13 Samadi et al. 2022	Adverse drug reactions recorded; treatment described as safe
14 Geohas et al. 2007	No significant adverse events in either group
16 2021	Product well tolerated
17 Sapp et al. 2026	Both well tolerated, no serious adverse events
18 Aksac et al. 2021	biotin prevented mucocutaneous adverse effects of isotretinoin
21 Schulpis et al. 2001	Skin lesions (seborrheic rash, alopecia) improved with biotin

6. What Remains Open: Limits of the Evidence

- Most controlled studies are small and short; large long-term studies are still pending.
- Whether the findings for hair, skin and nails are tied to a biotin deficiency is an open question.
- Many findings come from combination products and apply to the mixture, not to biotin alone.
- In multiple sclerosis, the results are inconsistent and only some of the endpoints were investigated.

- For most uses, controlled studies with clinical endpoints are still pending.

7. Bottom Line: What Is Established About Biotin (Vitamin B7)

Biotin is a water-soluble B vitamin that, as a coenzyme of several carboxylases, is involved in fat, carbohydrate and protein metabolism. In type 2 diabetes, biotin lowered total cholesterol in a meta-analysis; the effect on blood glucose is not yet confirmed. For hair, skin and nails, measured values improved in small studies mainly compared with baseline; the most informative placebo-controlled hair study found no difference. Many positive results come from combination products and therefore do not apply to biotin alone. In progressive multiple sclerosis, high-dose biotin showed benefit in an initial study, and the follow-up study did not confirm it. In signs of biotin deficiency, such as in pregnancy, biotin improved biotin-dependent metabolic values. Biotin was well tolerated; in high doses it can distort laboratory tests.

Frequently Asked Questions

How well has Biotin (Vitamin B7) been studied in humans?

We evaluated 123 human studies on Biotin (Vitamin B7), including 12 randomised controlled trials and 6 meta-analyses or systematic reviews; the controlled studies include 1,170 participants in total. As of 11 October 2026.

Does Biotin (Vitamin B7) help with biotin deficiency? What the studies show

We evaluated 3 controlled studies and 24 further studies; the controlled studies include 128 participants in total. Result of the controlled studies: mostly improved. Strength of evidence under our published rules: moderate. Sources: [10152022](#)

Does Biotin (Vitamin B7) help with multiple sclerosis? What the studies show

We evaluated 2 meta-analyses or systematic reviews, 2 controlled studies and 20 further studies; the controlled studies include 247 participants in total. Result of the controlled studies: mostly no difference. Strength of evidence under our published rules: low. Sources: [1579](#)

Does Biotin (Vitamin B7) help with hair loss? What the studies show

We evaluated one meta-analysis or systematic review, 2 controlled studies and 9 further studies; the controlled studies include 165 participants in total. Result of the controlled studies: mixed. Strength of evidence under our published rules: unclear. Sources: [481821](#)

Does Biotin (Vitamin B7) help with skin and skin inflammation? What the studies show

We evaluated 2 controlled studies and 5 further studies; the controlled studies include 165 participants in total. Result of the controlled studies: mostly no difference. Strength of evidence under our published rules: unclear. Sources: [8182152](#)

Does Biotin (Vitamin B7) help with glucose and lipid metabolism? What the studies show

We evaluated one meta-analysis or systematic review, one controlled study and 5 further studies; the controlled studies include 40 participants in total. Result of the controlled studies: mixed. Strength of evidence under our published rules: unclear. Sources: [231924](#)

Does Biotin (Vitamin B7) help with brittle nails? What the studies show

We evaluated one controlled study and 2 further studies; the controlled studies include 105 participants in total. Result of the controlled studies: mixed. Strength of evidence under our published rules: unclear. Sources: [897105](#)

What is biotin and what is it needed for in the body?

Biotin is a water-soluble vitamin from the group of B vitamins. In the body it acts as a coenzyme of several carboxylases that are involved in fat, carbohydrate and protein metabolism. Medically it is given in confirmed biotin deficiency and in congenital disorders of biotin metabolism.

Do human studies show an effect of biotin on hair loss?

A systematic review found three studies on biotin in hair loss; the most informative, placebo-controlled one showed no difference in hair growth ⁴. For biotin from a plant extract and for combination products with biotin, improvements were reported, but only compared with baseline or another hair product [81213](#).

How does biotin affect blood glucose and lipid metabolism?

In adults with type 2 diabetes, total cholesterol fell with biotin in a meta-analysis; the reduction in fasting blood glucose and triglycerides was not statistically confirmed, insulin remained unchanged ². For mixtures of biotin and chromium picolinate, two randomized controlled studies reported improvements in blood glucose and lipid values, but these findings apply to the mixture and not to biotin alone ⁶¹⁴.

What role does biotin play in multiple sclerosis?

In multiple sclerosis, high-dose biotin (MD1003) was tested against placebo in several randomized studies ⁷⁹²⁵. In a randomized study in progressive MS, disability regressed in some patients with high-dose biotin, and in no one with placebo ⁷. The follow-up study SPI2 did not confirm this benefit ²⁵. Another randomized controlled study showed no difference in visual acuity, nerve fiber layer thickness and quality of life in MS-related chronic vision loss ⁹. A systematic review with meta-analysis found no confirmed difference from placebo in disability and EDSS, but more often a faster walking time ¹.

Is biotin safe and what side effects are known?

Adverse events with high-dose biotin in multiple sclerosis occurred no more frequently than with placebo ¹⁷⁹; in children with asthma, biotin proved to be safe ¹¹. However, high-dose biotin distorted laboratory tests in some of those treated, and in no one with placebo ¹.

Sources

- 1 High-dose biotin for multiple sclerosis: A systematic review and meta-analyses of randomized controlled trials
Espiritu AI, Remalante-Rayco PPM · 2021
- 2 Influence of biotin intervention on glycemic control and lipid profile in patients with type 2 diabetes mellitus: A systematic review and meta-analysis
Zhang Y, Ding Y, Fan Y, Xu Y, Lu Y, Zhai L et al. · 2022
- 3 Effects of water-soluble vitamins on glycemic control and insulin resistance in adult type 2 diabetes: an umbrella review of meta-analyses
Chai Y, Chen C, Yin X, Wang X, Yu W, Pan H et al. · 2025
- 4 Biotin for Hair Loss: Teasing Out the Evidence
Yeilich A, Jenkins H, Holt S, Miller R · 2024
- 5 The effect of emerging nutraceutical interventions for clinical and biological outcomes in multiple sclerosis: A systematic review
Marx W, Hockey M, McGuinness AJ, Lane M, Christodoulou J, van der Mei I et al. · 2020
- 6 Combination of chromium and biotin improves coronary risk factors in hypercholesterolemic type 2 diabetes mellitus: a placebo-controlled, double-blind randomized clinical trial
Albarracín C, Fuqua B, Geohas J, Juturu V, Finch MR, Komorowski JR · 2007
- 7 MD1003 (high-dose biotin) for the treatment of progressive multiple sclerosis: A randomised, double-blind, placebo-controlled study
Tourbah A, Lebrun-Frenay C, Edan G, Clanet M, Papeix C, Vukusic S et al. · 2016
- 8 The Role of Sesbania grandiflora-Derived Biotin and Bambusa arundinacea-Derived Silica Extracts in Promoting Hair, Skin, and Nail Health: A Randomized, Double-Blind, Placebo-Controlled Clinical Study
Patel MN, Maheshvari J, Patel N · 2025
- 9 MD1003 (High-Dose Pharmaceutical-Grade Biotin) for the Treatment of Chronic Visual Loss Related to Optic Neuritis in Multiple Sclerosis: A Randomized, Double-Blind, Placebo-Controlled Study
Tourbah A, Gout O, Vighetto A, Deburghgraeve V, Pelletier J, Papeix C et al. · 2018
- 10 A Higher-Biotin Prenatal Multivitamin Formulation Improves Maternal and Cord Blood Biotin Biomarkers: A 24-Week Double-Blind, Randomized, Controlled Trial
Hanson I, Draicchio F, Sharafi M, Grier-Welch A, Vance A, Alamdari N et al. · 2026
- 11 L-Carnitine and Biotin as Adjunctive Therapies in Children with Moderate Persistent Asthma: A Randomized Placebo-Controlled Study
Khedr MO, El-Haggag SM, El-Razik AMA, elmeazawy RZ, Mostafa TM · 2026
- 12 A Comparative Study of Topical Procapil With Platelet-Rich Plasma Therapy Versus Topical Redensyl, Saw Palmetto, and Biotin With Platelet-Rich Plasma Therapy in the Treatment of Androgenetic Alopecia
Tr P, Ts R, K SK, Prasanna H · 2023
- 13 Efficacy of intramuscular injections of biotin and dexpantenol in the treatment of diffuse hair loss: A randomized, double-blind controlled study comparing two brands
Samadi A, Ketabi Y, Firooz R, Firooz A · 2022
- 14 Chromium picolinate and biotin combination reduces atherogenic index of plasma in patients with type 2 diabetes mellitus: a placebo-controlled, double-blinded, randomized clinical trial
Geohas J, Daly A, Juturu V, Finch M, Komorowski JR · 2007
- 15 Marginal biotin deficiency during normal pregnancy
Mock DM, Quirk JG, Mock NI · 2002
- 16 New Specialized Food Product "SPP1" in Combination With Iso-Calorie Diet Shows Better Efficacy Than the Diet Alone in Treatment of Non-alcoholic Steatohepatitis
· 2021
- 17 Absorption kinetics of vitamins and minerals from a novel nutritional product in physically active adults: a randomized, double-blind, placebo-controlled crossover trial
Sapp PA, Townsend JR, Kirby TO, Edwards CG, La Monica MB, Ziegenfuss TN et al. · 2026
- 18 Evaluation of biophysical skin parameters and hair changes in patients with acne vulgaris treated with isotretinoin, and the effect of biotin use on these parameters
Aksac SE, Bilgili SG, Yavuz GO, Yavuz IH, Aksac M, Karadag AS · 2021
- 19 Effects of biotin on lipids and other constituents of plasma of healthy men and women
Marshall MW, Kliman PG, Washington VA, Mackin JF, Weinland BT · 1980
- 20 Biotin supplementation affects lymphocyte carboxylases and plasma biotin in severe protein-energy malnutrition
Velázquez A, Terán M, Báez A, Gutiérrez J, Rodríguez R · 1995
- 21 Low serum biotinidase activity in children with valproic acid monotherapy
Schulpis KH, Karikas GA, Tjamouranis J, Regoutas S, Tsakiris S · 2001
- 22 Molecular Background and Disease Prevalence of Biotinidase Deficiency in a Polish Population-Data Based on the National Newborn Screening Programme
Jezela-Stanek A, Suchoń L, Sobczyńska-Tomaszewska A, Czarska K, Kuśmierska K, Taybert J et al. · 2022
- 23 Associations of dietary biotin intake on anxiety and depression: findings from a population-based prospective cohort study
Kong Y, Xia W, Wang T, Zhang D · 2025
- 24 Impairment of gut microbial biotin metabolism and host biotin status in severe obesity: effect of biotin and prebiotic supplementation on improved metabolism
Belda E, Voland L, Tremaroli V, Falony G, Adriouch S, Assmann KE et al. · 2022
- 25 Association Between Neurofilament Light Chain and Real-World Ambulatory Function in Progressive MS
Joseph GB, McCune M, Bove R, Abdelhak A, Cree BAC, Block VJ · 2026
- 26 Expanded Newborn Screening for Inborn Errors of Metabolism at a Single Center in Louisiana (2005-2024): Outcomes
Upadia J, Noh G, Crivelly K, Aziz E, Cunningham A, Andersson HC · 2025
- 27 Biotinidase biochemical and molecular analyses: Experience at a large reference laboratory
Sharma R, Kucera CR, Nery CR, Lacbawan FL, Salazar D, Tanpaiboon P · 2024
- 28 A retrospective study on biotinidase deficiency: analysis of the Eastern Anatolia region patient cohort
Yaralı O, Arslan S, Gündoğdu Öğütli ÖB, Guler MC, Akgül BN · 2024
- 29 Comprehensive analysis of genotypic and phenotypic characteristics of biotinidase deficiency patients in the eastern region of Türkiye
Çıki K, Alavanda C, Ceylan El, Tanyalçın T, Kilavuz S · 2024
- 30 Assessment of biotinidase activity changes over time in biotinidase deficient patients
Gülbağçe A, Muderrisoglu A, Güneş H, Erdoğan M, Dündar M, Kardaş F · 2026
- 31 Supplementary medication in multiple sclerosis: Real-world experience and potential interference with neurofilament light chain measurement
Pape K, Steffen F, Zipp F, Bittner S · 2020
- 32 High-dose biotin in progressive multiple sclerosis: A prospective study of 178 patients in routine clinical practice
Couloume L, Barbin L, Leray E, Wiertlewski S, Le Page E, Kerbrat A et al. · 2020
- 33 Hair Loss After Sleeve Gastrectomy and Effect of Biotin Supplements
Şen O, Türkçapar AG · 2021
- 34 Study of association between valproic acid levels, biotinidase activity and hair loss in Indian population: A cross-sectional comparative study
Kakunje A, Prabhu A, Karkal R, Pookoth KR, Kumar P, Gupta N et al. · 2025
- 35 Clinical, biochemical and genotypical characteristics in biotinidase deficiency
Akgun A, Sen A, Onal H · 2021
- 36 Associations of Biotin Levels in Serum and Follicular Fluid With ICSI Success: A Cross-Sectional Study From Iraq
Jaafar ZAA, Bakir NA, Salman DA · 2025
- 37 Biomarkers of treatment response in patients with progressive multiple sclerosis treated with high-dose pharmaceutical-grade biotin (MD1003)
Collongues N, Kuhle J, Tsagkas C, Lamy J, Meyer N, Barro C et al. · 2021
- 38 High dose biotin as treatment for progressive multiple sclerosis
Birnbau G, Stulc J · 2017
- 39 Hudson memorial lecture. Neonatal management of organic acidurias. Clinical update
Saudubray JM, Ogier H, Charpentier C, Depondt E, Coude FX, Munnich A et al. · 1984
- 40 Screening for Biotinidase deficiency in children with spinal cord demyelination
Mohammed M, El Agouza I, Zaitoun R, Ahmad K, Mohammad S, Sallam O et al. · 2026
- 41 Enzymatic Evolution and Longitudinal Recovery in Biotinidase Deficiency: Genotypic and Clinical Insights from the Follow-Up of a Newborn-Screened Cohort in Emilia-Romagna, Italy
Ortolano R, Menabò S, Candela E, Biasucci G, Bortolamedi E, Montanari G et al. · 2025
- 42 Exploring Secondary Biotinidase Deficiency and Biotin Supplementation in PMM2-CDG
Himmelreich N, Garbade SF, Okun JG, Hengst S, Geiger V, Barone R et al. · 2026
- 43 Short anagen syndrome: A case series and algorithm for diagnosis
Starace M, Gurioli C, Carpanese MA, Bruni F, Piraccini BM, Patrizi A et al. · 2021
- 44 The Clinical Evaluation of Serum WS Biotin, a Novel Encapsulated Form of D-Biotin With Improved Water Solubility, for Anti-Hair Shedding Applications. A Prospective Single-Arm, Nonrandomized, Pretest-Posttest Study
González Fernández D, Duchi S, Fernández Gómez L, León Sala T, Hajuj A, Molho D et al. · 2025
- 45 3-Hydroxypropionic acid and methylcitric acid are not reliable indicators of marginal biotin deficiency in humans
Mock DM, Henrich-Shell CL, Carnell N, Stumbo P, Mock NI · 2004
- 46 1H-31 P magnetic resonance spectroscopy: effect of biotin in multiple sclerosis
Guillevin C, Agius P, Naudin M, Herpe G, Ragot S, Maubeuge N et al. · 2019
- 47 High-dose pharmaceutical-grade biotin in patients with demyelinating neuropathies: a phase 2b open label, uncontrolled, pilot study
Créange A, Hutin E, Sedel F, Le Vigouroux L, Lefacheur JP · 2023
- 48 Neuroimaging Features of Biotinidase Deficiency
Biswas A, McNamara C, Gowda VK, Gala F, Sudhakar S, Sidpra J et al. · 2023
- 49 Sequence variants in the BTD underlying biotinidase deficiency in families of Pakistani origin
Moatter T, Ahmed S, Majid H, Jafri L, Bilal M, Najumuddin et al. · 2024
- 50 Lymphocyte propionyl-CoA carboxylase and its activation by biotin are sensitive indicators of marginal biotin deficiency in humans
Stratton SL, Bogusiewicz A, Mock MM, Mock NI, Wells AM, Mock DM · 2006
- 51 Biotinidase Deficiency in Children: Clinical Outcomes and Neuroimaging Correlations
Seilianian Toosi F, Saberifar M, Ahangari N, Hashemi N, Beiraghi Toosi M · 2026
- 52 Holocarboxylase Synthetase Deficiency: Clinical, Biochemical and Molecular Findings in Five Malaysian Patients Including a Newborn Presenting as Colloidion Baby
Ting SL, Yakob Y, Sani HA, Rethanavelu K, Ngu LH · 2025
- 53 The Clinical and Biochemical Impact of the Multivitamin Shortage on Neonatal Patients
Frankel R, Verscaj C, Tise CG, Cusmano-Ozog K · 2025
- 54 Recurrent "outsider" intronic variation in the SLC5A6 gene causes severe mixed axonal and demyelinating neuropathy, cyclic vomiting and optic atrophy in 3 families from Maghreb
Mansour-Hendili L, Gitiaux C, Harion M, Latouche C, Heron B, Stojkovic T et al. · 2024
- 55 Thiamine transporter-2 deficiency: outcome and treatment monitoring
Ortigoza-Escobar JD, Serrano M, Molero M, Oyazabal A, Rebollo M, Muchart J et al. · 2014
- 56 Profound Biotinidase Deficiency as a Treatable Cause of Pediatric Diffuse Leukoencephalopathy and Diffusion Restriction
Raha SJ, Saini AG, Attri SV, Bhatia V, Didwal G, Suthar R et al. · 2026
- 57 Atypical presentation of biotinidase deficiency: masquerading neuromyelitis optica spectrum disorder
Ali F, Mukhtiar K, Raza M, Ibrahim S · 2024
- 58 A novel SLC5A6 homozygous variant in a family with multivitamin-dependent neurometabolic disorder: Phenotype expansion and long-term follow-up
Montomoli M, Vetro A, Tubili F, Donati MA, Daniotti M, Pochiero F et al. · 2023
- 59 SLC5A6 Mutations in Axonal Sensorimotor Polyneuropathy Patients Concurrent With Sodium Dependent Multivitamin Transporter Deficiency and Improved Effects by Multivitamin Therapy
Pi BK, Lee AJ, Nam SH, Chung KW, Choi BO · 2025
- 60 Loss of taste responds to high-dose biotin treatment
Greenway FL, Ingram DK, Ravussin E, Hausmann M, Smith SR, Cox L et al. · 2011
- 61 Multiple Carboxylase Deficiency in an Infant Presenting With Severe Metabolic Acidosis and Sepsis-Like Features: A Case Report and Literature Review
Rehman T, Saeed R, Rauf MAA, Asif L, Junaid M, Khalid M et al. · 2026

- 62 A Short Indel is Causing Biotinidase Deficiency in a 4-Month-Old Boy from Jammu and Kashmir
Sharma R, Bhatti A, Singh D, Kapoor N, Bharathiraja S, Kaul S et al. · 2026
- 63 Precision diagnosis guides ketogenic diet therapy in inherited metabolic epilepsies: Concurrent non-ketotic hyperglycemia and biotinidase deficiency
Elbarky A, El Amrousy D, Mahmoud S, Lotfy A, Aboeisa M, Elballat KE et al. · 2026
- 64 Biotin-Induced Proximal Renal Tubular Acidosis in an Adolescent Female: Report of a Rare Case With Rechallenge Confirmation
Ahmad QT, Haseeb A, Ullah W, Samad M, Afridi A · 2026
- 65 Biotinidase Deficiency Induced Optic Neuropathy: A Case Report and Literature Review
Cao AA, Brown MM, Lee MS, Rashidi AM, Rashidi V · 2025
- 66 Expanded Clinical Phenotype and the Role of Untargeted Metabolomics Analysis in Confirming the Diagnosis of Sodium-Dependent Multivitamin Transporter Deficiency
Walimbe AS, Waskow E, Mackay L, Miller M, Gijavanekar C, Difalco CR et al. · 2025
- 67 Hepatoerythropoietic Porphyria with Coexisting BDT and $CNGB1$ Genetic Mutations: A First Case Report
Kaya Ç, Heydari A, Kızılay O, Kızılay O, Sezgin G · 2025
- 68 Prenatal Brain Abnormalities in Sodium-Dependent Multivitamin Transporter Deficiency
Ogawa E, Kosaki K, Takenouchi T · 2025
- 69 Clinical diagnosis, treatment, and genetic analysis of adolescent onset holocarboxylase synthetase deficiency and cobalamin C deficiency: A case report and literature review
Ren Y, Dang H, Fu Y, Liu C, Li J, Cai J · 2025
- 70 Biotinidase deficiency deterioration in the second decade, presenting as treatable cerebellar ataxia and encephalopathy masquerading as demyelination
Gowda VK, Badrinath AC, Srinivasan VM, Verma MP, Nayyer A · 2025
- 71 Biotin-Thiamine-Responsive Basal Ganglia Disease: A Case Report
Tse J, Abu-Qamar A, Youssef O, Pejka SL · 2025
- 72 High-Dose Biotin Supplementation Exacerbates Chronic Migraine: A Case Report
Sanosi AA · 2025
- 73 A new case of sodium-dependent multivitamin transporter defect occurring as a life-threatening condition responsive to early vitamin supplementation and literature review
Van Vyve FX, Mercier N, Papadopoulos J, Heijmans C, Dessy H, Monestier O et al. · 2024
- 74 Optic Neuropathy and Myelopathy in a Teenager With Biotinidase Deficiency
Chodnicki KD, Aksamit AJ, Gavrilova RH, Farnsworth PJ, McClelland CM · 2024
- 75 Dramatic Clinical Improvement With Biotin Mega-Dose Therapy in a Neonate With Holocarboxylase Synthetase Deficiency
Kim SW, Lee HJ, Choi N, Kim EK, Ko JM · 2024
- 76 Ophthalmic manifestations of biotinidase deficiency: report of a case and review of literature
Abdi F, Parvin S, Zare Hosseiniabadi V, Kachuei M, Gordiz A, Hemmati S et al. · 2024
- 77 Biotinidase Deficiency in the Second Decade with Atypical Neuroimaging Findings
Gowda VK, Avaragollapuravarga Mathada A, Srinivasan VM, Vamyanmane DK · 2023
- 78 Developing of Biotin-Thiamine Responsive Basal Ganglia Disease after Accidental Ingestion of Ethyl Alcohol: A Case Report
Aldosari AN, Arisha A, Ibrahim A, Gongli M · 2023
- 79 Two novel BTD mutations causing profound biotinidase deficiency in a Chinese patient
Geng J, Sun Y, Zhao Y, Xiong W, Zhong M, Zhang Y et al. · 2021
- 80 Tardive Reactivation of Progressive Multiple Sclerosis During Treatment with Biotin
Demas A, Cochlin JP, Hardy C, Vaschalde Y, Bourre B, Labauge P · 2020
- 81 A Case Report of Solitary Sclerosis: This is Really Multiple Sclerosis
Lebrun C, Cohen M, Mondot L, Ayrygnac X, Labauge P · 2017
- 82 Untargeted Metabolomic and Lipidomic Profiling Reveals Distinct Biochemical Patterns in Treated Biotinidase Deficiency
Ünlü Torlak E, Koç Yekedüz M, Bülbül YE, Kara İS, Erdoğan Kablan S, Eylem CC et al. · 2026
- 83 Characterization of a Novel BTD Hypomorphic Variant in a Patient with Complex Neurodevelopmental Delay: Resolving Actionable Metabolic Vulnerabilities Beyond Borderline Plasma Biochemistry
Toledo-Pacheco C, Montero-Hernández M, López-Garrido MP, de Diego-Boguñá C, Sánchez-Sánchez F · 2026
- 84 Biotinidase Deficiency: Report of a Tunisian Case With Neuromyelitis Optica-Like Presentation and Review of the Literature
Zioudi A, Benrhouma H, Jamoussi M, Ben Younes T, Miladi Z, Klau H et al. · 2025
- 85 Biotinidase deficiency presenting with status epilepticus and cerebellar infarction in an infant: a case report
Shrestha B, Karn V, Regmi N, Aryal S, Khadka A, Neupane S · 2025
- 86 Evaluation of clinical, laboratory, and molecular genetic features of patients with biotinidase deficiency
Yılmaz B, Ceylan AC, Gündüz M, Ünal Uzun Ö, Küçükcongar Yavaş A, Bilginer Gürbüz B et al. · 2024
- 87 A comprehensive investigation of biochemical status in patients with telogen effluvium: Analysis of Hb, ferritin, vitamin B12, vitamin D, thyroid function tests, zinc, copper, biotin, and selenium levels
Durusu Turkoglu IN, Turkoglu AK, Soyulu S, Gencer G, Duman R · 2024
- 88 Elevated C5-hydroxy acylcarnitine in an infant girl as a result of holocarboxylase synthetase deficiency
Nelson AT, Jones PM, Cao J · 2024
- 89 Alterations in optical coherence tomography and optical coherence tomography angiography findings in children with partial biotinidase deficiency
Karataş G, Çakır A, Çakar NE, Çepni Çakır D, Karataş ME, Akpunar O et al. · 2024
- 90 What Dietary Vitamins and Minerals Might Be Protective against Parkinson's Disease?
Alizadeh M, Kheirouri S, Keramati M · 2023
- 91 Novel SLC5A6 mutations lead to B lymphocyte maturation defects with metabolic abnormality rescuable by biotin replenishment
Hsieh CH, Lee J, Sung HH, Huang YF, Ding YS, Li CY et al. · 2023
- 92 A Retrospective Evaluation of the Laboratory Findings of Dermatology Patients Whose Biotin Levels Were Checked
Güder H, Eker P · 2023
- 93 A mild case of sodium-dependent multivitamin transporter (SMVT) deficiency illustrating the importance of treatment response in variant classification
Hauth I, Waterham HR, Wanders RJA, van der Crabben SN, van Karnebeek CDM · 2022
- 94 Relapses in Patients Treated with High-Dose Biotin for Progressive Multiple Sclerosis
Mathais S, Moisset X, Pereira B, Taithe F, Ciron J, Labauge P et al. · 2021
- 95 Relapses During High-Dose Biotin Treatment in Progressive Multiple Sclerosis: a Case-Crossover and Propensity Score-Adjusted Prospective Cohort
Branger P, Parienti JJ, Derache N, Kassiss N, Assouad R, Maillart E et al. · 2020
- 96 Biotin Deficiency in Telogen Effluvium: Fact or Fiction?
Abdel Rahman SH, Mohammed Salem R, Hassan Sabry J · 2020
- 97 Transungual Delivery, Anti-Inflammatory Activity, and In Vivo Assessment of a Cyclodextrin Polypseudotaxanes Nail Lacquer
Fernández-Campos F, Navarro F, Corrales A, Picas J, Pena E, González J et al. · 2020
- 98 Nutritional Adjuvants in Diabetes: Biotin and Chromium Picolinate as Emerging Therapeutic Agents
Mehta M, Saini A, Mehta DK, Das R · 2026
- 99 The Potential Neuroprotective Role of Biotin in Neurorehabilitation: A Review
Gunay E, Gezmen-Karadag M · 2026
- 100 Alpha-Lipoic Acid and Biotin in Neurodegenerative Diseases: Convergent Mechanistic Insights from Preclinical Models to Clinical Perspectives
Aguilera-Méndez A, Aguilera-Manuel K, Saavedra-Molina A, Ríos-Chávez P, Villafaña S, Nieto-Aguilar R et al. · 2026
- 101 Holocarboxylase synthetase deficiency: pathogenesis, clinical features, diagnosis, treatment, and research prospects
ZeZhao J, Ablimit A · 2026
- 102 [Hair loss during treatment of obesity: current state of the problem and potential solutions]
Troshina EA, Suplotova LA, Dzgoeva FK, Ruyatkina LA, Gadzhigorieva AG, Gromova OA et al. · 2026
- 103 Biotinidase Deficiency
Woodliff B, Benjamin R, Simpson B · 2025
- 104 Primary Progressive Multiple Sclerosis: New Therapeutic Approaches
Rajabi M, Shafaeibajestan S, Asadpour S, Alvari G, Taei N, Kohkalanli M et al. · 2025
- 105 Nail Supplements: When, How, and Why?
Zaraa I, Richert B · 2025
- 106 Comparative bioavailability of vitamins in human foods sourced from animals and plants
Chungchunlam SMS, Moughan PJ · 2024
- 107 Biotin Homeostasis and Human Disorders: Recent Findings and Perspectives
Karachaliou CE, Livanlou E · 2024
- 108 Nutritional Supplements for Skin Health-A Review of What Should Be Chosen and Why
Januszewski J, Forma A, Zembala J, Flieger M, Tyczyńska M, Dring JC et al. · 2023
- 109 Valproate-associated hair abnormalities: Pathophysiology and management strategies
Prahraj SK, Munoli RN, Udupa ST, Vaidyanathan S · 2022
- 110 [Progress and prospects of metabolic therapy in multiple sclerosis]
Bisaga GN, Mikhailenko AA, Barsukov IN · 2019
- 111 Use of Vitamins and Dietary Supplements by Patients With Multiple Sclerosis: A Review
Evans E, Piccio L, Cross AH · 2018
- 112 The Evidence for Dietary Interventions and Nutritional Supplements as Treatment Options in Multiple Sclerosis: a Review
Mische LJ, Mowry EM · 2018
- 113 Actual Insights into Treatable Inborn Errors of Metabolism Causing Epilepsy
Mastrangelo M · 2018
- 114 Treatment of progressive multiple sclerosis: Challenges and promising perspectives
Maillart E · 2018
- 115 Pharmacokinetics and pharmacodynamics of MD1003 (high-dose biotin) in the treatment of progressive multiple sclerosis
Peyro Saint Paul L, Debruyne D, Bernard D, Mock DM, Defer GL · 2016
- 116 Targeting demyelination and virtual hypoxia with high-dose biotin as a treatment for progressive multiple sclerosis
Sedel F, Bernard D, Mock DM, Tourbah A · 2016
- 117 Intersections of pathways involving biotin and iron relative to therapeutic mechanisms for progressive multiple sclerosis
Heidker RM, Emerson MR, LeVine SM · 2016
- 118 Biotin-Thiamine-Responsive Basal Ganglia Disease
Tabarki B, Al-Hashem A, Ortigoza-Escobar JD, Alsharhan H, Alfadhel M · 1993

International Health Research

Research, put in context. Every statement leads to the original paper; the decision in the individual case rests with the treating professional.
ih-research.com/en/knowledge/biotin/ · Last updated 2026-10-17